

**Table 5.9 Broadleaf Weed Control in Bluegrass, Tall Fescue, Perennial Ryegrass, and Common Bermudagrass**

Weed	Classification	Response to Herbicides					Preferred Time to Treat
		2,4-D + dicamba	2,4-D + MCPP	2,4-D + Dicamba+ MCPP	2,4-D + Dicamba+ Triclopyr	2,4-D + Dicamba + Penoxsulam	
<b>Bedstraw</b>	<sup>1</sup> A	<sup>2</sup> S	I	R-I	S	I-S	Spring
<b>Bindweed - Field Hedge</b>	P	S	I-S	S	S	S	Spring
<b>Bittercress</b>	WA	S	S	S	S	S	Fall
<b>Black Medic</b>	SA & P	S	I	S	S	S	Spring & Fall
<b>Buttercup</b>	WA & P	S	S	S	S	S	Spring & Fall
<b>Buttonweed - Virginia</b>	P	R-I	R	R-I	I-S	I-S	Spring Sequentially
<b>Carpetweed</b>	SA	S	S	S	S	S	Spring
<b>Carrot - Wild</b>	B	S	S	S	S	S	Fall
<b>Catsear Dandelion</b>	P	S	S	S	S	S	Fall
<b>Chickweed - Common</b>	WA	S	S	S	S	S	Spring & Fall
<b>Mouseear</b>	WA & P	S	I-S	S	S	S	Spring & Fall
<b>Chicory</b>	P	S	S	S	S	S	Fall
<b>Cinquefoil - Common</b>	A	S	S	S	S	S	Spring
<b>Clover - Crimson</b>	SA	S	S	S	S	S	Spring
<b>Hop</b>	SA	S	S	S	S	S	Spring
<b>White</b>	P	S	S	S	S	S	Spring & Fall
<b>Daisy - Oxeye</b>	P	I	I	I	I-S	S	Spring & Fall
<b>Dandelion</b>	P	S	S	S	S	S	Spring & Fall
<b>Dichondra</b>	P	I-S	I-S	I-S	I-S	I-S	Spring & Fall
<b>Dock</b>	P	I-S	I	I-S	I-S	I-S	Spring
<b>Dogfennel</b>	P	I	R-I	I	I-S	I-S	Spring & Fall
<b>Garlic - Wild</b>	P	I	I	I	I	I	Fall
<b>Geranium - Carolina</b>	WA	S	S	S	S	S	Spring & Fall
<b>Ground ivy</b>	P	I	I	I	I-S	I-S	Spring
<b>Hawkweed</b>	P	I	I	I	I-S	I-S	Fall
<b>Healall</b>	P	S	I-S	S	S	S	Fall
<b>Henbit</b>	WA	I-S	I	S	S	S	Fall
<b>Honeysuckle</b>	P	S	I-S	S	S	S	Spring & Summer
<b>Horsenettle</b>	P	I	R-I	I	I	I	Spring
<b>Horseweed</b>	WA & SA	I-S	I-S	I-S	S	I-S	Spring & Fall
<b>Knapweed</b>	B	I	I	I	I-S	I-S	Fall
<b>Knawel</b>	WA	I	I	I	I-S	I-S	Fall
<b>Knotweed</b>	SA	I	I	I	I-S	I-S	Spring & Summer
<b>Lambsquarters</b>	SA	S	S	S	S	S	Spring & Summer
<b>Lespedeza</b>	SA	I	I	I	I-S	I	Spring

**Table 5.9 Broadleaf Weed Control in Bluegrass, Tall Fescue, Perennial Ryegrass, and Common Bermudagrass (cont.)**

Weed	Classification	Response to Herbicides					Preferred Time to Treat
		2,4-D + dicamba	2,4-D + MCPP	2,4-D + Dicamba+ MCPP	2,4-D + Dicamba+ Triclopyr	2,4-D + Dicamba + Penoxsulam	
<b>Mallow</b>	SA	I-S	I	I-S	I-S	I-S	Spring & Summer
<b>Mugwort</b>	P	R-I	R-I	R-I	I	I	Spring
<b>Mustards</b>	WA & B	I-S	I	S	S	S	Fall
<b>Onion - Wild</b>	P	I	I	I	I	I	Fall
<b>Pennycress</b>	A	S	S	S	S	S	Fall
<b>Pepperweed</b>	WA	S	S	S	S	S	Fall
<b>Pigweed</b>	SA	S	S	S	S	S	Spring & Summer
<b>Plantains</b>	P	S	S	S	S	S	Spring & Fall
<b>Poison ivy</b>	P	I-S	I	I-S	I-S	I-S	Summer & Fall
<b>Poorjoe</b>	A	I	I	I	I-S	I-S	Spring & Summer
<b>Spurge - Prostrate</b>	SA	R-I	R-I	R-I	R-I	I-S	Summer
<b>Spotted</b>	SA	R-I	R-I	R-I	R-I	I-S	Summer
<b>Purslane</b>	SA	S	I-S	S	S	S	Spring & Summer
<b>Red Sorrel</b>	P	I	I	I	I-S	I-S	Fall
<b>Shepherd's Purse</b>	WA	S	S	S	S	S	Fall
<b>Smartweed</b>	SA	I-S	I	I-S	I-S	I-S	Spring & Summer
<b>Sowthistle</b>	WA	S	S	S	S	S	Fall
<b>Speedwell - Corn</b>	WA	R	R	R	I	--	Spring
<b>Persian</b>	WA	R	R	R	I	--	Spring
<b>Star-of-Bethlehem</b>	P	R	R	R	R	--	Spring
<b>Strawberry - Wild</b>	P	I	R-I	I	I	I	Fall
<b>Teasel - Common</b>	B	I-S	I-S	I-S	S	I-S	Spring
<b>Cutleaf</b>	B	I-S	I-S	I-S	S	I-S	Spring
<b>Thistle - Bull</b>	B	I-S	I-S	I-S	I-S	R-I	Fall
<b>Canada</b>	P	I	R-I	I	I	R-I	Fall
<b>Musk</b>	B	S	S	S	S	S	Spring
<b>Plumless</b>	B & P	S	S	S	S	I-S	Spring
<b>Violet</b>	P	R-I	R	R-I	I-S	I-S	Spring Sequentially
<b>Wood sorrel - Creeping</b>	WA & P	R-I	R-I	R-I	I-S	--	Spring
<b>Yellow</b>	A	I	I	I	I-S	--	Spring
<b>Yarrow</b>	P	I	I	I	S	--	Fall
<b>Yellow Rocket</b>	B & P	S	I-S	S	S	I-S	Fall

<sup>1</sup>A = annual (summer or winter); B = biennial; P = perennial; SA = summer annual; WA = winter annual.

<sup>2</sup>R = resistant in most instances, poor control usually less than 50%; I = intermediate tolerance, good control at times with high rates, may require multiple treatments; S = weed susceptibility, often good control 70% or higher.